

6. (Reiterated) The antigen presenting vesicle according to claim 13, wherein said antigen presenting cell is derived from a B-lymphocyte, a Langerhans cell, a macrophage or a dendritic cell.

9. (Three Times Amended) An antigen presenting vesicle having a membrane and a major histocompatibility complex (MHC) class I protein, obtained by the step of:

recovering a membrane-enriched fraction obtained by differential centrifugation of membrane-containing fractions of cell culture supernatants or lysates of antigen presenting cells.

*None done*

11. (Amended) An antigen presenting vesicle having a membrane and a major histocompatibility complex class I molecule, obtained by the step of:

recovering a 70,000 x g pellet obtained by differential centrifugation of membrane-containing fractions of cell culture media or lysates of antigen presenting cells containing MHC class I.

*done*

12. (Amended) A purified antigen presenting vesicle having a membrane and a major histocompatibility complex class I molecule, obtained by the step of:

recovering a fraction having a buoyant density of 1.10 to 1.22 g/ml from a 70,000 x g pellet obtained by differential centrifugation of membrane-containing fractions of cell culture supernatants or lysates of antigen presenting cells containing MHC class I.

*done*

13. (Twice Amended) An antigen presenting vesicle free from its natural surroundings, comprising:

a membrane and a major histocompatibility complex (MHC) class I protein or a functional derivative or fragment thereof, wherein said antigen presenting vesicle is obtainable from an antigen presenting cell.